State of Michigan 2020Q4 Aggregate TAP Report





Michigan Department of Health and Human Services
Surveillance for Healthcare-Associated and Resistant Pathogens (SHARP) Unit

The Michigan Department of Health and Human Services (MDHHS) Surveillance for Healthcare-Associated and Resistant Pathogens (SHARP) Unit began including the new targeted assessment for prevention (TAP) reports in the 2014 annual statewide aggregate report. Aggregate reports are available for acute care hospitals in each emergency preparedness region and critical access hospitals in all regions.

This report shows modules and locations where the State of Michigan either needs to focus additional prevention efforts or is excelling in infection prevention. **Table 1** presents a cumulative attributable difference (CAD) determined using the 2020 HHS target standardized infection ratios (SIRs) for each module, using the NHSN 2015 Baselines, which is modeled after the data included in the CDC National and State Annual HAI Report. Numbers with "Need to Prevent" next to them show how many infections the state needs to prevent quarterly in order to reach the 2020 HHS target SIR. Numbers with "Prevented" next to them show the number of infections prevented beyond what was expected for the state according to the 2020 HHS target SIR. Corresponding SIRs for each module and location type are provided as well.

NHSN Module ¹	Hospital Type	Number of Facilities ²	Location ³	SIR ⁴	Significant (Y/N) ⁵	CAD ⁶	Prevented or Need to Prevent
CAUTI	Acute	88	All	0.850	Υ	21.330	Need to Prevent
		72	ICU	0.863	N	12.445	Need to Prevent
		88	WARD+	0.835	N	8.886	Need to Prevent
	Critical Access	15	All	0.670	N	-0.119	Prevented
			ICU				
		15	WARD+	0.724	N	-0.036	Prevented
CLABSI	Acute	88	All	0.857	N	18.896	Need to Prevent
		72	ICU	1.083	N	25.520	Need to Prevent
		88	WARD+	0.716	Υ	-3.048	Prevented
		18	NICU	0.396	Υ	-1.051	Prevented
	Critical Access	16	All			-0.176	Prevented
			ICU				
		16	WARD+			-0.151	Prevented
			NICU				
CDI	Acute	90	Facility-wide	0.595	Y	-73.067	Prevented
	Critical Access	16	Facility-wide	0.450	N	-1.111	Prevented
MRSA Bac	Acute	90	Facility-wide	0.851	N	28.446	Need to Prevent
	Critical Access	16	Facility-wide			-0.189	Prevented
SSI COLO	All	81		1.073	N	25.047	Need to Prevent
SSI HYST	All	79		0.914	N	3.510	Need to Prevent
SSI HPRO	All	69		1.271	N	12.135	Need to Prevent
SSI KPRO	All	68		0.717	N	0.235	Need to Prevent

¹CAUTI, catheter-associated urinary tract infection; CLABSI, central line-associated bloodstream infection; CDI, *Clostridium difficile* infection LabID; MRSA Bac, methicillin-resistant *Staphylococcus aureus* bloodstream infection LabID; SSI COLO, admission/readmission colon surgical site infection; SSI HYST, admission/readmission abdominal hysterectomy surgical site infection; SSI HPRO, admission/readmission hip arthroplasty surgical site infection; SSI KPRO, admission/readmission knee arthroplasty surgical site infection.

²Note: facilities in which an SIR could not be calculated with a CAD of 0 were excluded from this table. Data for locations with less than 5 facilities reporting have been suppressed.

³All includes all units for which in-plan data are reported; ICU includes all critical care units; WARD+ includes all WARD, WARD_ONC, SCA, STEP, or OTHER units; NICU includes all neonatal critical care units; Facility-wide includes all inpatient units.

⁴SIR: Standardized Infection Ratio: Ratio of observed events compared to the number of predicted events, accounting for unit type or other variables. An SIR of 1 can be interpreted as having the same number of events as predicted. An SIR that is between 0 and 1 represents fewer events than predicted, while an SIR of greater than 1 represents more events than predicted. SIRs were calculated using the 2015 NHSN Baselines. ⁵Significant (Y/N). A Y indicates that, based on the p-value and 95% Confidence Interval (CI), the SIR is statistically significantly different than 1. An N indicates that, based on the p-value and 95% CI, the SIR is not statistically significantly different than 1 (expected).

⁶CAD=Cumulative Attributable Difference. The number of infections that the state either needs to prevent to meet the 2020 HHS target or has prevented beyond the 2020 HHS target. 2020 HHS HAI Target SIRs: CAUTI = 0.75, CLABSI = 0.50, CDI = 0.70, MRSA bacteremia = 0.50, SSI = 0.70.

In addition to the NHSN modules reported above, other module data are reported to the MDHHS SHARP Unit. However, there are no corresponding HHS Target SIRs for these HAIs and therefore CAD values are not calculated. SIR values using the NHSN 2015 Baseline for these HAIs are shown in **Table 2** below.

Table 2. 2020Q4 Standardized Infection Ratios for Other HAIs							
NHSN Module ¹	Hospital Type	Number of Facilities ²	Location ³	SIR ⁴	Significant (Y/N) ⁵		
Total VAE	Acute	76	All	1.374	Υ		
		58	ICU	1.430	Υ		
		58	WARD+	0.920	N		
	Critical Access	11	All				
			ICU				
		9	WARD+				
IVAC	Acute	76	All	1.461	Υ		
		58	ICU	1.520	Υ		
		58	WARD+	0.957	N		
MBI-CLABSI	Acute	88	All	1.258	N		
		72	ICU	1.354	N		
		88	WARD+	1.245	N		

¹Total VAE, total ventilator-associated events; IVAC Plus, infection-related ventilator-associated complication; MBI-CLABSI, mucosal barrier injury central line-associated bloodstream infection.

²Note: facilities for which the number of predicted infections are 0 have been excluded. Data for locations with less than 5 facilities reporting have been suppressed.

³All includes all units for which in-plan data are reported; ICU includes all critical care units; WARD+ includes all WARD, WARD_ONC, SCA, STEP, or OTHER units; NICU includes all neonatal critical care units; Facility-wide includes all inpatient units.

⁴SIR: Standardized Infection Ratio: Ratio of observed events compared to the number of predicted events, accounting for unit type or other variables. An SIR of 1 can be interpreted as having the same number of events as predicted. An SIR that is between 0 and 1 represents fewer events than predicted, while an SIR of greater than 1 represents more events than predicted. SIRs were calculated using the 2015 NHSN Baselines.

⁵Significant (Y/N). A Y indicates that, based on the p-value and 95% Confidence Interval (CI), the SIR is statistically significantly different than 1. An N indicates that, based on the p-value and 95% CI, the SIR is not statistically significantly different than 1 (expected).

Standardized utilization ratios (SURs) is a risk-adjusted comparison of the amount of indwelling device use to the amount expected to be used for each facility using the NHSN 2015 Baselines. There are currently no national or state targets for SURs. The SUR data reported to the MDHHS SHARP unit is included in **Table 3** below.

Table 3. 2020Q4 Standardized Utilization Ratios for Indwelling Device Use							
Indwelling Device Type	Hospital Type	Number of Facilities ¹	Location ²	SUR ³	Significant (Y/N) ⁴		
Central Lines	Acute	88	All	0.875	Υ		
		72	ICU	0.985	Υ		
		88	WARD+	0.985	Υ		
		18	NICU	0.868	Υ		
	Critical Access	16	All	1.167	N		
			ICU				
		16	WARD+	1.881	Υ		
Urinary Catheters	Acute	88	All	0.787	Υ		
		72	ICU	0.978	Υ		
		88	WARD+	0.978	Υ		
	Critical Access	15	All	0.835	Υ		
			ICU				
		15	WARD+	0.928	N		
Ventilators	Acute	76	All	0.919	N		
		58	ICU	1.168	Υ		
		76	WARD+	1.168	Υ		
	Critical Access	11	All	1.061	Υ		
			ICU				
		9	WARD+	0.509	Υ		

¹Note: facilities for which the number of predicted device days are 0 have been excluded. Data for locations with less than 5 facilities reporting have been suppressed.

Please contact Elli Stier (<u>RayE7@michigan.gov</u>) with questions, comments, or suggestions. All aggregate reports are posted at <u>www.michigan.gov/hai</u>.

²All includes all units for which in-plan data are reported; ICU includes all critical care units; WARD+ includes all WARD, WARD_ONC, SCA, STEP, or OTHER units; NICU includes all neonatal critical care units; Facility-wide includes all inpatient units.

³SUR: Standardized Utilization Ratio: Ratio of observed device days compared to the number of predicted device days, accounting for unit type or other variables. An SUR of 1 can be interpreted as having the same number of device days as predicted. An SUR that is between 0 and 1 represents fewer device days than predicted, while an SUR of greater than 1 represents more device days than predicted. SURs were calculated using the 2015 Baselines.

⁴Significant (Y/N). A Y indicates that, based on the p-value and 95% Confidence Interval (CI), the SUR is statistically significantly different than 1. An N indicates that, based on the p-value and 95% CI, the SUR is not statistically significantly different than 1 (expected).